

Optimizing Hospital Length of Stay, Quality Perspective

Altman M, Vanpée M, Cnattingius S, Norman M. Moderately preterm infants and determinants of length of hospital stay. Arch Dis Child Fetal Neonatal Ed. 2009 Nov;94(6):F414-8. doi: 10.1136/adc.2008.153668. Epub 2009 May 21. PubMed PMID: 19465411.

Eichenwald EC, Zupancic JA, Mao WY, Richardson DK, McCormick MC, Escobar GJ. Variation in diagnosis of apnea in moderately preterm infants predicts length of stay. Pediatrics. 2011 Jan;127(1):e53-8. doi: 10.1542/peds.2010-0495. Epub 2010 Dec 27. PubMed PMID: 21187315.

Eichenwald EC, Blackwell M, Lloyd JS, Tran T, Wilker RE, Richardson DK. Inter-neonatal intensive care unit variation in discharge timing: influence of apnea and feeding management. Pediatrics. 2001 Oct;108(4):928-33. PubMed PMID: 11581446.

Escobar GJ, Joffe S, Gardner MN, Armstrong MA, Folck BF, Carpenter DM. Rehospitalization in the first two weeks after discharge from the neonatal intensive care unit. Pediatrics. 1999 Jul;104(1):e2. PubMed PMID: 10390288.

Goyal N, Zubizarreta JR, Small DS, Lorch SA. Length of stay and readmission among late preterm infants: an instrumental variable approach. Hosp Pediatr. 2013 Jan;3(1):7-15. PubMed PMID: 24319830; PubMed Central PMCID: PMC3967867.

Grossman MR, Berkowitz AK, Osborn RR, Xu Y, Esserman DA, Shapiro ED, Bizzarro MJ. An Initiative to Improve the Quality of Care of Infants With Neonatal Abstinence Syndrome. Pediatrics. 2017 Jun;139(6). pii: e20163360. doi: 10.1542/peds.2016-3360. Epub 2017 May 18. PubMed PMID: 28562267; PubMed Central PMCID: PMC5470506.

Ingram J, Blair PS, Powell JE, Manns S, Burden H, et al. Preparing for Home: a before-and-after study to investigate the effects of a neonatal discharge package aimed at increasing parental knowledge, understanding and confidence in caring for their preterm infant before and after discharge from hospital. Southampton (UK): NIHR Journals Library; 2016 Mar. PubMed PMID: 26985528.

Jadcherla SR, Dail J, Malkar MB, McClead R, Kelleher K, Nelin L. Impact of Process Optimization and Quality Improvement Measures on Neonatal Feeding Outcomes at an All-Referral Neonatal Intensive Care Unit. JPEN J Parenter Enteral Nutr. 2016 Jul;40(5):646-55. doi: 10.1177/0148607115571667. Epub 2015 Mar 2. PubMed PMID: 25733339.

Karagiannidou S, Zaoutis T, Maniadas N, Papaevangelou V, Kourlaba G. Attributable length of stay and cost for pediatric and neonatal central line-associated bloodstream infections in Greece. J Infect Public Health. 2019 May - Jun;12(3):372-379. doi: 10.1016/j.jiph.2018.12.004. Epub 2019 Jan 4. PubMed PMID: 30616938.

Lapcharoensap W, Gage SC, Kan P, Profit J, Shaw GM, Gould JB, Stevenson DK, O'Brodovich H, Lee HC. Hospital variation and risk factors for bronchopulmonary dysplasia in a population-based cohort. JAMA Pediatr. 2015 Feb;169(2):e143676. doi: 10.1001/jamapediatrics.2014.3676. Epub 2015 Feb 2. PubMed PMID: 25642906.

Lee HC, Bennett MV, Crockett M, Crowe R, Gwiazdowski SG, et al. Comparison of Collaborative Versus Single-Site Quality Improvement to Reduce NICU Length of Stay. Pediatrics. 2018 Jul;142(1). pii: e20171395. doi: 10.1542/peds.2017-1395. Epub 2018 Jun 13. PubMed PMID: 29899043.

Maier RF, Blondel B, Piedvache A, Misselwitz B, Petrou S, et al. Duration and Time Trends in Hospital Stay for Very Preterm Infants Differ Across European Regions. *Pediatr Crit Care Med*. 2018 Dec;19(12):1153-1161. doi: 10.1097/PCC.0000000000001756. PubMed PMID: 30334907; PubMed Central PMCID: PMC6282674.

Mansfield SA, Ryshen G, Dail J, Gossard M, McClead R, Aldrink JH. Use of quality improvement (QI) methodology to decrease length of stay (LOS) for newborns with uncomplicated gastroschisis. *J Pediatr Surg*. 2018 Aug;53(8):1578-1583. doi: 10.1016/j.jpedsurg.2017.11.061. Epub 2017 Dec 8. PubMed PMID: 29291893.

Olsen SL, Park ND, Tracy K, Younger D, Anderson B. Implementing Standardized Feeding Guidelines, Challenges, and Results. *Neonatal Netw*. 2018 Jul;37(4):218-223. doi: 10.1891/0730-0832.37.4.218. Epub 2018 Jul 1. PubMed PMID: 30567919.

Proffit J, Zupancic JA, McCormick MC, Richardson DK, Escobar GJ, Tucker J, Tarnow-Mordi W, Parry G. Moderately premature infants at Kaiser Permanente Medical Care Program in California are discharged home earlier than their peers in Massachusetts and the United Kingdom. *Arch Dis Child Fetal Neonatal Ed*. 2006 Jul;91(4):F245-50. Epub 2006 Jan 31. PubMed PMID: 16449257; PubMed Central PMCID: PMC2672723.

Rodrigues C, Severo M, Zeitlin J, Barros H. The Type of Feeding at Discharge of Very Preterm Infants: Neonatal Intensive Care Units Policies and Practices Make a Difference. *Breastfeed Med*. 2017 Dec 6. doi: 10.1089/bfm.2017.0135. [Epub ahead of print] PubMed PMID: 29211544.

Settle M, Francis K. Does the Infant-Driven Feeding Method Positively Impact Preterm Infant Feeding Outcomes? *Adv Neonatal Care*. 2019 Feb;19(1):51-55. doi: 10.1097/ANC.0000000000000577. PubMed PMID: 30672812.

Silber JH, Lorch SA, Rosenbaum PR, Medoff-Cooper B, Bakewell-Sachs S, Millman A, Mi L, Even-Shoshan O, Escobar GJ. Time to send the preemie home? Additional maturity at discharge and subsequent health care costs and outcomes. *Health Serv Res*. 2009 Apr;44(2 Pt 1):444-63. doi: 10.1111/j.1475-6773.2008.00938.x. Epub 2008 Dec 31. PubMed PMID: 19207592; PubMed Central PMCID: PMC2677048.

Welch CD, Check J, O'Shea TM. Improving care collaboration for NICU patients to decrease length of stay and readmission rate. *BMJ Open Qual*. 2017 Oct 21;6(2):e000130. doi: 10.1136/bmjopen-2017-000130. eCollection 2017. PubMed PMID: 29450288; PubMed Central PMCID: PMC5699126.