

Improving Nutrition of the Extremely Preterm Infant

Abdelmaaboud M, Mohammed A. Early versus late minimal enteral feeding in weeks preterm growth-restricted neonates with abnormal antenatal Doppler studies. *J Matern Fetal Neonatal Med* 2015;(just-accepted):1–20.

Ågren J, Sjörs G, Sedin G. Transepidermal water loss in infants born at 24 and 25 weeks of gestation. *Acta Paediatr* 2007;87(11):1185–90.

Balasubramanian H, Nanavati RN, Kabra NS. Effect of two different doses of parenteral amino acid supplementation on postnatal growth of very low birth weight neonates, a randomized controlled trial. *Indian Pediatr* 2013;50(12):1131–6.

Benavente-Fernández I, Synnes A, Grunau RE, et al. Association of Socioeconomic Status and Brain Injury With Neurodevelopmental Outcomes of Very Preterm Children. *JAMA Netw Open* 2019;2(5):e192914.

Blanco CL, Falck A, Green BK, Cornell JE, Gong AK. Metabolic responses to early and high protein supplementation in a randomized trial evaluating the prevention of hyperkalemia in extremely low birth weight infants. *J Pediatr* 2008;153(4):535–40.

Blanco CL, Gong AK, Schoolfield J, et al. Impact of early and high amino acid supplementation on ELBW infants at 2 years. *J Pediatr Gastroenterol Nutr* 2012;54(5):601–7

Burattini I, Bellagamba MP, Spagnoli C, et al. Targeting 2.5 versus 4 g/kg/day of amino acids for extremely low birth weight infants: a randomized clinical trial. *J Pediatr* 2013;163(5):1278–82.e1.

Cakir B, Hellström W, Tomita Y, et al. IGF1, serum glucose, and retinopathy of prematurity in extremely preterm infants. *JCI Insight* [Internet] 2020;5(19). Available from: <http://dx.doi.org/10.1172/jci.insight.140363>

Castro M, Asbury M, Shama S, et al. Energy and Fat Intake for Preterm Infants Fed Donor Milk Is Significantly Impacted by Enteral Feeding Method. *JPEN J Parenter Enteral Nutr* 2018

Chacham S, Pasi R, Chegondi M, Ahmad N, Mohanty SB. Metabolic Bone Disease in Premature Neonates: An Unmet Challenge. *J Clin Res Pediatr Endocrinol* 2020;12(4):332–9.

Chan B, Lian A, Baer V, et al. An Evaluation to Establish the Acceptable Serum Triglyceride Levels in Neonates Receiving Intravenous Fat Emulsion Infusion in a Multicenter Retrospective Study. *Am J Perinatol* [Internet] 2020; Available from: <http://dx.doi.org/10.1055/s-0040-1705174>

Clyman R, Wickremasinghe A, Jhaveri N, et al. Enteral feeding during indomethacin and ibuprofen treatment of a patent ductus arteriosus. *J Pediatr* 2013;163(2):406.

Collins CT, Makrides M, McPhee AJ. Docosahexaenoic Acid and Bronchopulmonary Dysplasia in Preterm Infants. *N Engl J Med* 2017;376(13):1245–55.

Dorling J, Abbott J, Berrington J, et al. Controlled Trial of Two Incremental Milk-Feeding Rates in Preterm Infants. *N Engl J Med* 2019;381(15):1434–43.

Faienza MF, D'Amato E, Natale MP, et al. Metabolic Bone Disease of Prematurity: Diagnosis and Management. *Front Pediatr* 2019;7:143.

Fenton TR, Singhal N, Baynton RD, Akierman A. Serum triglycerides of breast milk-fed very-low-birth-weight infants. *Nutr Clin Pract* 1997;12(1):26–9.

Francis J, Unger S, Bando N, et al. Postdischarge Feeding of Very-low-birth-weight Infants: Adherence to Nutrition Guidelines. *J Pediatr Gastroenterol Nutr* 2018;67(3):401–8.

Garg P, Pinotti R, Lal CV, Salas AA. Transfusion-associated necrotizing enterocolitis in preterm infants: An updated meta-analysis of observational data. *J Perinat Med* 2017

Geftter WB, Epstein DM, Anday EK, Dalinka MK. Rickets presenting as multiple fractures in premature infants on hyperalimentation. *Radiology* 1982;142(2):371–4.

Gokçe İK, Oguz SS. Late onset hyponatremia in preterm newborns: is the sodium content of human milk fortifier insufficient? *J Matern Fetal Neonatal Med* 2020;33(7):1197–202.

Havranek T, Johanboeke P, Madramootoo C, Carver JD. Umbilical artery catheters do not affect intestinal blood flow responses to minimal enteral feedings. *J Perinatol* 2007;27(6):375.

Hay S, Zupancic JAF, Flannery DD, Kirpalani H, Dukhovny D. Should we believe in transfusion-associated enterocolitis? Applying a GRADE to the literature. *Semin Perinatol* 2017;41(1):80–91.

Jensen EA, Zhang H, Feng R, et al. Individualising care in severe bronchopulmonary dysplasia: a series of N-of-1 trials comparing transpyloric and gastric feeding. *Arch Dis Child Fetal Neonatal Ed* 2020;105(4):399–404.

Kantorowska A, Wei JC, Cohen RS, Lawrence RA, Gould JB, Lee HC. Impact of Donor Milk Availability on Breast Milk Use and Necrotizing Enterocolitis Rates. *Pediatrics* 2016;137(3):e20153123.

Koletzko, Poindexter, and Uauy. *Nutritional Care of Preterm Infants: Scientific Basis and Practical Guidelines*. S. Karger AG, 2014

Koletzko B, Poindexter B, Uauy R (eds): *Nutritional Care of Preterm Infants: Scientific Basis and Practical Guidelines*. *World Rev Nutr Diet*. Basel, Karger, 2014, vol 110, pp 177–189

Konnikova Y, et al. *PLOS One*. 2015;10:e0132924.

Leenders EKSM, de Waard M, van Goudoever JB. Low- versus High-Dose and Early versus Late Parenteral Amino-Acid Administration in Very-Low-Birth-Weight Infants: A Systematic Review and Meta-Analysis. *Neonatology* 2018;113(3):187–205.

Marc I, Piedboeuf B, Lacaze-Masmonteil T, et al. Effect of Maternal Docosahexaenoic Acid Supplementation on Bronchopulmonary Dysplasia-Free Survival in Breastfed Preterm Infants: A Randomized Clinical Trial. *JAMA* 2020;324(2):157–67.

Martini S, Aceti A, Galletti S, Beghetti I, Faldella G, Corvaglia L. To Feed or Not to Feed: A Critical Overview of Enteral Feeding Management and Gastrointestinal Complications in Preterm Neonates with a Patent Ductus Arteriosus. *Nutrients* [Internet] 2019;12(1).

- Niinikoski H, Stoll B, Guan X, Kansagra K, Lambert BD, Stephens J, Hartmann B, Holst JJ, Burrin DG. Onset of small intestinal atrophy is associated with reduced intestinal blood flow in TPN-fed neonatal piglets. *J Nutr*. 2004 Jun;134(6):1467-74. doi: 10.1093/jn/134.6.1467. PMID: 15173413.
- Oddie SJ, Young L, McGuire W. Slow advancement of enteral feed volumes to prevent necrotising enterocolitis in very low birth weight infants. *Cochrane Database Syst Rev* 2017;8:2147483647.
- Parker MG, Gupta M, Melvin P, et al. Racial and Ethnic Disparities in the Use of Mother's Milk Feeding for Very Low Birth Weight Infants in Massachusetts. *J Pediatr* 2019;204:134–41.e1.
- Parker MG, Greenberg LT, Edwards EM, Ehret D, Belfort MB, Horbar JD. National Trends in the Provision of Human Milk at Hospital Discharge Among Very Low-Birth-Weight Infants. *JAMA Pediatr* [Internet] 2019
- Patole SK, de Klerk N. Impact of standardised feeding regimens on incidence of neonatal necrotising enterocolitis: A systematic review and meta-analysis of observational studies. *Arch Dis Child Fetal Neonatal Ed* 2005;90(2):51.
- Poindexter BB, Martin CR. Impact of Nutrition on Bronchopulmonary Dysplasia. *Clin Perinatol*. 2015 Dec;42(4):797-806. doi: 10.1016/j.clp.2015.08.007. Epub 2015 Oct 2. PMID: 26593079.
- Radmacher PG, Adamkin DH. Fortification of human milk for preterm infants. *Semin Fetal Neonatal Med* 2017;22(1):30; 2 O'Connor DL, Kiss A, Tomlinson C, et al. Nutrient enrichment of human milk with human and bovine milk-based fortifiers for infants born weighing <1250 g: a randomized clinical trial. *Am J Clin Nutr* 2018;108(1):108–16.
- Rochow N, Fusch G, Choi A, et al. Target fortification of breast milk with fat, protein, and carbohydrates for preterm infants. *J Pediatr* 2013;163(4):1001.
- Rochow N, Landau-Crangle E, So HY, et al. Z-score differences based on cross-sectional growth charts do not reflect the growth rate of very low birth weight infants. *PLoS One* 2019;14(5):e0216048.
- Sangild PT. Gut responses to enteral nutrition in preterm infants and animals. *Exp Biol Med (Maywood)*. 2006 Dec;231(11):1695-711. doi: 10.1177/153537020623101106. PMID: 17138756.
- Segar DE, Segar EK, Harshman LA, Dagle JM, Carlson SJ, Segar JL. Physiological Approach to Sodium Supplementation in Preterm Infants. *Am J Perinatol* 2018;35(10):994.
- Simon L, Théveniaut C, Flamant C, Frondas-Chauty A, Darmaun D, Rozé J-C. In Preterm Infants, Length Growth below Expected Growth during Hospital Stay Predicts Poor Neurodevelopment at 2 Years. *Neonatology* 2018;114(2):135–41.
- Sisk PM, Lovelady CA, Dillard RG, Gruber KJ, O'Shea TM. Maternal and infant characteristics associated with human milk feeding in very low birth weight infants. *J Hum Lact* 2009;25(4):412–9.
- Strydom K, Van Niekerk E, Dhansay MA. Factors affecting body composition in preterm infants: Assessment techniques and nutritional interventions. *Pediatr Neonatol* 2019;60(2):121–8.
- Sunehag AL, Haymond MW, Schanler RJ, Reeds PJ, Bier DM. Gluconeogenesis in very low birth weight infants receiving total parenteral nutrition. *Diabetes* 1999;48(4):791.

Sung SI, Ahn SY, Seo HJ, et al. Insensible Water Loss during the First Week of Life of Extremely Low Birth Weight Infants Less than 25 Gestational Weeks under High Humidification. Neonatal Medicine 2013;20(1):51–7.

Thureen PJ, Melara D, Fennessey PV, Hay WW Jr. Effect of low versus high intravenous amino acid intake on very low birth weight infants in the early neonatal period. Pediatr Res 2003;53(1):24–32.

Uthaya S, Liu X, Babalis D, et al. Nutritional Evaluation and Optimisation in Neonates (NEON) trial of amino acid regimen and intravenous lipid composition in preterm parenteral nutrition: a randomised double-blind controlled trial. Efficacy and Mechanism Evaluation 2016;3(2):1.

Wu PY, Hodgman JE. Insensible water loss in preterm infants: changes with postnatal development and non-ionizing radiant energy. Pediatrics 1974;54(6):704–12.

Yeo KT, Kong JY, Sasi A, Tan K, Lai NM, Schindler T. Stopping enteral feeds for prevention of transfusion-associated necrotising enterocolitis in preterm infants. Cochrane Database Syst Rev [Internet] 2019;2019(10).

<https://www.bapm.org/>