

Addressing Comorbidities Through Implementation of an Exclusive Human Milk Diet (EHMD)

References

Abrams SA, Schanler RJ, Lee ML, Rechtman DJ. Greater mortality and morbidity in extremely preterm infants fed a diet containing cow milk protein products. *Breastfeed Med*. 2014 Jul-Aug;9(6):281-5. doi: 10.1089/bfm.2014.0024. Epub 2014 May 27. PMID: 24867268; PMCID: PMC4074755.

Agostoni C, Buonocore G, Carnielli VP, De Curtis M, Darmaun D, et al; ESPGHAN Committee on Nutrition. Enteral nutrient supply for preterm infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2010 Jan;50(1):85-91. doi: 10.1097/MPG.0b013e3181adaee0. PMID: 19881390.

Assad M, Elliott MJ, Abraham JH. Decreased cost and improved feeding tolerance in VLBW infants fed an exclusive human milk diet. *J Perinatol*. 2016 Mar;36(3):216-20. doi: 10.1038/jp.2015.168. Epub 2015 Nov 12. PMID: 26562370.

Cristofalo EA, Schanler RJ, Blanco CL, Sullivan S, Trawoeger R, et al. Randomized trial of exclusive human milk versus preterm formula diets in extremely premature infants. *J Pediatr*. 2013 Dec;163(6):1592-1595.e1. doi: 10.1016/j.jpeds.2013.07.011. Epub 2013 Aug 20. PMID: 23968744.

Fleig L, Hagan J, Lee ML, Abrams SA, Hawthorne KM, Hair AB. Growth outcomes of small for gestational age preterm infants before and after implementation of an exclusive human milk-based diet. *J Perinatol*. 2021 Aug;41(8):1859-1864. doi: 10.1038/s41372-021-01082-x. Epub 2021 May 19. PMID: 34012050; PMCID: PMC8342303.

Ganapathy V, Hay JW, Kim JH. Costs of necrotizing enterocolitis and cost-effectiveness of exclusively human milk-based products in feeding extremely premature infants. *Breastfeed Med*. 2012 Feb;7(1):29-37. doi: 10.1089/bfm.2011.0002. Epub 2011 Jun 30. PMID: 21718117.

Ghandehari H, Lee ML, Rechtman DJ; H2MF Study Group. An exclusive human milk-based diet in extremely premature infants reduces the probability of remaining on total parenteral nutrition: a reanalysis of the data. *BMC Res Notes*. 2012 Apr 25;5:188. doi: 10.1186/1756-0500-5-188. PMID: 22534258; PMCID: PMC3527141.

Hair AB, Peluso AM, Hawthorne KM, Perez J, Smith DP, et al. Beyond Necrotizing Enterocolitis Prevention: Improving Outcomes with an Exclusive Human Milk-Based Diet. *Breastfeed Med*. 2016 Mar;11(2):70-4. doi: 10.1089/bfm.2015.0134. Epub 2016 Jan 20. Erratum in: *Breastfeed Med*. 2017 Dec;12(10):663. PMID: 26789484; PMCID: PMC4782036.

Hair AB, Hawthorne KM, Chetta KE, Abrams SA. Human milk feeding supports adequate growth in infants \leq 1250 grams birth weight. *BMC Res Notes*. 2013 Nov 13;6:459. doi: 10.1186/1756-0500-6-459. PMID: 24220185; PMCID: PMC3879715.

Hair AB, Blanco CL, Moreira AG, Hawthorne KM, Lee ML, et al. Randomized trial of human milk cream as a supplement to standard fortification of an exclusive human milk-based diet in infants 750-1250 g birth weight. *J Pediatr*. 2014 Nov;165(5):915-20. doi: 10.1016/j.jpeds.2014.07.005. Epub 2014 Aug 15. PMID: 25130571.



Huston RK, Markell AM, McCulley EA, Gardiner SK, Sweeney SL. Improving Growth for Infants ≤ 1250 Grams Receiving an Exclusive Human Milk Diet. *Nutr Clin Pract*. 2018 Oct;33(5):671-678. doi: 10.1002/ncp.10054. Epub 2018 Feb 16. PMID: 29451716.

Huston RK, Markell AM, McCulley EA, et al. Decreasing Necrotizing Enterocolitis and Gastrointestinal Bleeding in the Neonatal Intensive Care Unit: The Role of Donor Human Milk and Exclusive Human Milk Diets in Infants ≤ 1500 g Birth Weight. *ICAN: Infant, Child, & Adolescent Nutrition*. 2014;6(2):86-93. doi:[10.1177/1941406413519267](https://doi.org/10.1177/1941406413519267)

Koletzko B, Poindexter B, Uauy R. Recommended nutrient intake levels for stable, fully enterally fed very low birth weight infants. *World Rev Nutr Diet*. 2014;110:297-9. doi: 10.1159/000360195. Epub 2014 Apr 11. PMID: 24751638.

O'Connor DL, Kiss A, Tomlinson C, Bando N, Bayliss A, et al; OptiMoM Feeding Group. Nutrient enrichment of human milk with human and bovine milk-based fortifiers for infants born weighing < 1250 g: a randomized clinical trial. *Am J Clin Nutr*. 2018 Jul 1;108(1):108-116. doi: 10.1093/ajcn/nqy067. Erratum in: *Am J Clin Nutr*. 2019 Aug 1;110(2):529. Erratum in: *Am J Clin Nutr*. 2020 May 1;111(5):1112. PMID: 29878061.

Parra-Llorca A, Gormaz M, Alcántara C, Cernada M, Nuñez-Ramiro A, et al. Preterm Gut Microbiome Depending on Feeding Type: Significance of Donor Human Milk. *Front Microbiol*. 2018 Jun 27;9:1376. doi: 10.3389/fmicb.2018.01376. PMID: 29997594; PMCID: PMC6030370.

Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 2012 Mar;129(3):e827-41. doi: 10.1542/peds.2011-3552. Epub 2012 Feb 27. PMID: 22371471.

Sullivan S, Schanler RJ, Kim JH, Patel AL, Trawöger R, et al. An exclusively human milk-based diet is associated with a lower rate of necrotizing enterocolitis than a diet of human milk and bovine milk-based products. *J Pediatr*. 2010 Apr;156(4):562-7.e1. doi: 10.1016/j.jpeds.2009.10.040. Epub 2009 Dec 29. PMID: 20036378.

Visuthranukul C, Abrams SA, Hawthorne KM, Hagan JL, Hair AB. Premature small for gestational age infants fed an exclusive human milk-based diet achieve catch-up growth without metabolic consequences at 2 years of age. *Arch Dis Child Fetal Neonatal Ed*. 2019 May;104(3):F242-F247. doi: 10.1136/archdischild-2017-314547. Epub 2018 Nov 13. PMID: 30425116; PMCID: PMC6764250.